Diagnostic criteria in Pai syndrome: results of a case series and a literature review

Anne Morice, Eva Galliani, Jeanne Amiel, Martin Rachwalski, Cecilia Neiva, C. Thauvin-Robinet, Marie-Paule Vasquez, Arnaud Picard, Natacha Kadlub

► To cite this version:

Anne Morice, Eva Galliani, Jeanne Amiel, Martin Rachwalski, Cecilia Neiva, et al.. Diagnostic criteria in Pai syndrome: results of a case series and a literature review. *International Journal of Oral and Maxillofacial Surgery*, 2019, 48 (3), pp.Pages 283-290. hal-01948492

HAL Id: hal-01948492

<https://u-bourgogne.hal.science/hal-01948492>

Submitted on 21 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Distributed under a Creative Commons Attribution - NonCommercial| 4.0 International License

[ARTICLE TYPE – CLINICAL PAPER, CRANIOFACIAL ANOMALIES]

Diagnostic criteria in Pai syndrome: results of a case series and a literature review

**Anne Morice^{1,2}, Eva Galliani¹, Jeanne Amiel^{2,3,4}, Martin Rachwalski^{1,4,5}, Cecilia
Neiva¹, Christel Thauvin-Robinet⁶, Marie-Paule Vazquez^{1,2}, Arnaud Picard^{1,2},
Natacha Kadlub^{1,2}**

*¹Department of Maxillofacial and Plastic Surgery, Rare Diseases Reference Centre
Coordinator for Clefts and Facial Malformations, Hôpital Universitaire Necker-
Enfants Malades, Paris, France*

²Université Paris Descartes-Sorbonne Paris Cité, Paris, France

*³Department of Medical Genetics, Hôpital Universitaire Necker-Enfants Malades,
Paris, France*

*⁴Imagine Institute of Genetic Diseases, Université Paris Descartes-Sorbonne Paris
Cité, Paris, France*

*⁵Department of Paediatric Neurosurgery, National Reference Centre for Craniofacial
Malformations, Hôpital Universitaire Necker-Enfants Malades, Paris, France*

*⁶Department of Genetic Diseases, Reference Centre for Developmental Anomalies
and Malformative Syndromes, Centre Hospitalier Universitaire, Dijon, France*

Abstract

Pai syndrome was originally described as the association of a midline cleft lip, midline facial polyps, and lipoma of the central nervous system. However, only a few patients present with the full triad, and most exhibit a wide spectrum of phenotypic variability. The aim of this study was to phenotypically delineate Pai syndrome and to propose new criteria to facilitate a clinical diagnosis in the future. The study cohort consisted of seven case patients and an additional 60 cases diagnosed with Pai syndrome identified in a literature review. Only 23 of 67 patients presented the full triad as historically described by Pai et al. (1987). A congenital facial midline skin mass was always encountered, particularly affecting the nasal structures (60/67). A midline facial cleft was reported in 45 of 67 patients and a pericallosal lipoma in 42 of 67 patients. The proposed definition of Pai syndrome is the association of (1) a congenital nasal or mediofrontal skin mass or a mid-anterior alveolar process polyp as a mandatory criterion, and at least one of the following criteria: (2) midline cleft lip and/or midline alveolar cleft, and/or (3) a pericallosal lipoma or interhemispheric lipoma in the case of corpus callosum dysgenesis.

Key words: Pai syndrome, midline cleft lip, skin tag, pericallosal lipoma, frontonasal dysplasia

Introduction

Pai syndrome is a rare congenital craniofacial malformation involving the midline cranial and facial structures. According to the literature, Pai syndrome is

phenotypically highly variable and overlaps with frontonasal dysplasia spectrum in some cases¹⁻³. Most of the cases of Pai syndrome described in the literature do not meet all criteria as initially defined by Pai et al., i.e., (1) a midline cleft lip, (2) midline nasal and facial polyps, and (3) lipoma of the central nervous system⁴. Less stringent diagnostic criteria have been proposed, including the presence of at least (1) a midline cleft lip, and (2) one hamartomatous nasal polyp and/or mid-anterior alveolar process polyp⁵, or the association of a congenital nasal polyp and one of the three following criteria: (1) midline cleft lip, (2) mid-anterior alveolar process congenital polyp, or (3) a pericallosal lipoma³.

Most cases are sporadic, and although an autosomal dominant inheritance has been suggested, the aetiology remains unknown^{2,6}. Establishing homogeneous study cohorts in the future may help to elucidate the aetiology of Pai syndrome. Thus, there is the need for a consensus definition of Pai syndrome.

This article describes a series of seven additional patients diagnosed with Pai syndrome and provides a review of cases reported in the literature according to the three current clinical diagnostic criteria. The aim of this study was first to describe the phenotype spectrum of reported cases of Pai syndrome and to determine whether these cases are consistent with the existing definitions of Pai syndrome. The second aim was to propose more appropriate diagnostic criteria and systematic investigations, based on these findings.

Patients and methods

Patients

A retrospective study was conducted in the Rare Diseases Reference Centre for Cleft Lip and Palate and Facial Malformations at Hôpital Trousseau and Hôpital Universitaire Necker-Enfants Malades, Paris, France, covering the years 2008 to 2018. Using the CEMARA database (Banque Nationale de Données Maladies Rares), seven patients diagnosed with Pai syndrome were identified. These patients presented at least two criteria of the triad as defined by Pai et al.⁴, with a midline facial congenital skin mass as a mandatory criterion. The following information was collected from a chart review: sex, birth parameters, location of the congenital facial skin mass, description of the facial cleft and additional abnormalities, neuropsychological development, and results of genetic investigations. Cerebral magnetic resonance imaging (MRI) scans were reviewed to determine the type of pericallosal lipoma and any additional central nervous system malformations.

Literature review

The review of the literature was conducted by screening the PubMed database using the terms “Pai syndrome” and/or “midline lip cleft” and/or “facial skin mass” and/or “cutaneous polyps” and/or “lipoma of the corpus callosum”. This analysis included case reports and series of Pai syndrome, as well as literature reviews published since 1965.

Results

Case series (Table 1, Figs. 1–3)

[Table 1 here]

[Figures 1–3 here]

Seven patients were included in this study, five female and two male, with a median age of 1 month (range 0.25–5 months) at initial examination and a median follow-up of 9 years (range 0.5–19 years). For all patients, family history was unremarkable, and the patient's parents were not consanguineous. Patients 5 and 7 each had a non-affected dizygotic twin. Cleft lip was diagnosed prenatally in two patients (patients 2 and 7).

Among the seven patients, four presented with the full triad (patients 1–4) as described by Pai et al.⁴, whereas three patients presented two criteria (patients 5–7; Table 1; Fig. 1). A midline nasal skin mass was present in all cases, a midline facial cleft was observed in six of seven cases, and a pericallosal lipoma was seen in five of seven cases (Fig. 3 [Au?1]). Multiple congenital skin tags were present in three of seven patients (Table 1; Fig. 2 [Au?2]).

Ophthalmological abnormalities were diagnosed in two patients, including a hyperpigmented spot of the iris and a limbal dermoid. Intracranial abnormalities were also detected, including partial agenesis of the corpus callosum, short corpus callosum, choroid cyst, and agenesis of the olfactory bulbs. Hypertelorism, bifid nose, and down-slanting palpebral fissures were noted in patient 3 (Fig. 1). Two patients presented a bifid midline frenum of the upper lip (Fig. 2 [Au?2]). Neuropsychological development was considered in the normal range for all patients.

Literature review

Fifty articles matched the research criteria, including 46 case reports and four case series (from two to seven patients per series). Among these case reports and series, nine included a review of the literature^{3,5,7–13}.

In total, 60 cases of Pai syndrome have been described to date. Nineteen patients met the three clinical criteria as defined by Pai et al.⁴ (**Supplementary Material**, Table S1). Thirty-six patients presented two of the three criteria: 14 patients had at least one nasal skin mass and a midline cleft of the upper lip, 15 patients had at least one congenital nasal skin mass and a pericallosal lipoma, and seven patients presented with a paramedian cleft lip or midline alveolar cleft without cleft lip, associated with midline facial polyps and/or a pericallosal lipoma (**Supplementary Material**, Table S2). Forty-two cases featured additional abnormalities, with multiple congenital facial skin masses (40/60), hypertelorism (19/60), and ophthalmological anomalies (19/60) being the most frequently observed (see **Supplementary Material**, Tables S1 and S2). Five patients presented one criterion only (i.e., nasal congenital skin mass).

Discussion

The phenotypic spectrum of the reported cases of Pai syndrome is heterogeneous², as are the diagnostic criteria^{3–5}, which was confirmed by the present review. This study first described the phenotype of seven additional cases of Pai syndrome and then compared them with 60 cases of Pai syndrome reported in the literature to date, in order to determine whether these cases are consistent with the existing definitions of Pai syndrome. The second aim was to critically analyse the three definitions of Pai

syndrome and, on the basis of the results, to propose new diagnostic criteria as well as systematic investigations.

Analysis of the clinical spectrum of Pai syndrome

Congenital facial skin masses

Almost all cases reported in the literature, as well as the cases in the series presented herein, harboured a facial midline hamartoma (57/60 cases in the literature and 6/7 cases in this series). Castori et al. have previously noted that this is the most frequently encountered feature⁵. These congenital skin masses affected the nasal structures in 53 of 60 cases reported in the literature, particularly the nostrils, and this was also seen in the present case series. In a few cases, examination of the skin masses revealed a lipoma or a myolipoma^{14–19} or a dermoid cyst⁶, as in two patients in the case series (see Table 1).

Midline cleft lip

Midline cleft lip is rare and represents 0.43% to 0.73% of cases of cleft lip and palate^{9,20}. In the literature, midline clefts were reported in 39 of 60 patients with Pai syndrome (65% [Au23]), with a wide range of severity exhibited (**Supplementary Material**, Tables S1 and S2). In the case series, six of seven patients had a midline cleft lip, and one patient presented with a paramedian cleft lip, a median notch of the upper lip, and a midline alveolar cleft (see Table 1). In the literature, three patients

with Pai syndrome presented with uni- or bilateral paramedian lip clefts^{2,21,22}. In four cases, a midline alveolar cleft without cleft lip was reported³.

Pericallosal lipoma

Intracranial lipomas represent less than 0.1% of intracranial anomalies. They are mostly located in the midline^{22,23}. Anterior lipomas of the corpus callosum are mostly of tubulonodular type and frequently associated with hypoplasia or agenesis of the corpus callosum, frontal encephalocele, frontal lobe anomalies, and ocular anomalies. Posterior lipomas are typically described as curvilinear; they are thin and typically lie on the splenium. They are rarely associated with corpus callosum and/or encephalic anomalies^{23,24}. A pericallosal lipoma or interhemispheric lipoma was identified in 37 of 60 patients with Pai syndrome; however, cerebral imaging was available for only 49 of the 60 patients. In the case series presented, five patients had a corpus callosum lipoma (tubulonodular in four cases and curvilinear type in the other). One patient with a tubulonodular pericallosal lipoma also presented partial agenesis of the corpus callosum.

Intracranial lipomas are rarely symptomatic²³, and in Pai syndrome, neuropsychological development is usually normal³, as confirmed in the case series. However, neuropsychological development is not well studied in Pai syndrome, and several conditions reported in Pai syndrome are likely to cause neuropsychological impairments. These include associated chromosomal abnormalities such as translocation q28;q11.2²⁵, intracranial calcifications^{9,26}, and corpus callosum dysgenesis^{9,27} or agenesis^{2,5,19}. Patients with Pai syndrome with corpus callosum dysgenesis²⁷, hypoplasia or agenesis^{2,19}, or other cerebral malformations^{6,28} often

present with an interhemispheric lipoma (**Supplementary Material**, Table S2). Thus, systematic cerebral imaging is recommended when a congenital midline facial skin mass and/or midline facial cleft is encountered, as there may be associated cerebral malformations^{6,28} [Au?4]. Cerebral imaging studies are imperative, as a pericallosal lipoma may become symptomatic at a later age, with the risk of developing epilepsy, which has been reported to occur in up to 50% of patients²⁹ [Au?4].

Additional features

The most frequent additional features encountered in the literature review were multiple congenital facial skin masses (66%), hypertelorism (32%), and ophthalmological anomalies (32%). In the case series, three of seven patients presented with multiple congenital facial skin masses. Two patients showed ophthalmological anomalies: a hyperpigmented spot of the iris and an ocular dermoid. The clinical spectrum of ophthalmological anomalies described in the literature is wide (**Supplementary Material**, Tables S1 and S2). However, Castori et al. suggested that they could be underestimated⁵. Thus, systematic and complete ophthalmological examination is justified when encountering Pai syndrome.

Ear anomalies are rare and these included pre-auricular skin tags and/or ear dysplasia in five of 60 patients^{21,30}. In the case series, one patient had three left pre-auricular skin tags and bilateral, symmetrical external ear dysplasia in addition to an ocular dermoid. The presence of ear dysplasia and epibulbar dermoids raises the possibility that there might be an overlap with Goldenhar/oculo-auriculo-vertebral (OAV) spectrum and oculoauriculofrontonasal syndrome, which is associated with features of both frontonasal dysplasia and OAV spectrum.

Defining Pai syndrome

To date, three definitions of Pai syndrome have been described^{3–5}. Pai et al. postulated that three criteria, i.e., a midline cleft lip, midline facial and nasal skin polyps, and lipoma of the corpus callosum, must be present⁴, whereas according to Castori et al.⁵ and Lederer et al.³, at least two criteria are sufficient: one or more hamartomatous nasal polyp(s) plus a midline facial cleft (midline cleft lip with or without cleft alveolus) and/or mid-anterior alveolar process congenital polyp (Castori et al.⁵) or the presence of a congenital nasal polyp plus one of the following three features: midline cleft lip (with or without cleft alveolus), mid-anterior alveolar process congenital polyp, or pericallosal lipoma (Lederer et al.³).

In all three definitions, a midline congenital skin polyp must be present. According to Pai et al.⁴, the congenital skin polyp may be located not only in the nasal region but also in the midline facial regions, whereas Castori et al.⁵ and Lederer et al.³ restricted this to the nasal structure only. Contrary to Lederer et al.³, a midline cleft lip must be present to define Pai syndrome in the two other definitions. Pericallosal lipoma is an obligatory criterion according to Pai et al.⁴, a possible associated feature in the definition of Lederer et al.³, and excluded from the diagnostic criteria of Castori et al.⁵.

In the literature review, less than a third of the cases [Au?5] (19/60) met the criteria defined by Pai et al.⁴, 34 of 60 cases met the Castori et al. criteria⁵, and 50 of 60 cases corresponded to the definition of Lederer et al.³. In the case series of seven patients, four presented the triad as defined by Pai et al.⁴ and six met the criteria described by Castori et al.⁵ and Lederer et al.³. Patient 6 did not meet the existing

definitions, as she presented with a congenital dermoid cyst of the nasal tip and not a hamartomatous polyp, in addition to a pericallosal lipoma without a midline cleft lip. Several cases diagnosed with Pai syndrome reported in the literature did not present with a hamartomatous polyp but rather with a congenital lipoma or a myolipoma^{14–19} or a dermoid cyst⁶. The pathological diagnosis is not described systematically in previous reports; thus, it is possible that the number of cases without a true hamartomatous polyp may have been underestimated [Au?4]. In addition, these congenital skin masses diagnosed as dermoid cysts or lipoma correspond to congenital and non-progressive lesions of developmental origin that may occur in the midline facial structures. In accordance with the existing definitions, it is considered that a midline congenital skin mass should be a mandatory criterion for defining Pai syndrome, as it was present in all of the cases in the literature review, occurring in the frontal and/or nasal regions as in the definition of Pai et al.⁴. In addition, it is proposed that the possible location be broadened to the mid-anterior alveolar process and its possible pathological diagnoses to dermoid cysts and hamartomatous-like lesions, such as lipoma or myolipoma.

In fact, the initial definition of Pai et al. is relatively restrictive. However, although this congenital malformation affecting the craniofacial midline structures can be expressed in various degrees, it can nevertheless be explained by similar pathophysiological processes [Au?4]. From a pathophysiological point of view, two theories may explain the occurrence of a median cleft of the upper lip: the ‘fusion theory’ and Veau’s theory of mesodermal penetration. According to the fusion theory, the cleft of the upper lip is due to a lack of fusion of the maxillary processes with the frontonasal process during foetal face development³¹. In Veau’s theory, the median cleft of the upper lip results from abnormal mesenchymal penetration in the epithelial

wall of the primary palate during the sixth week of gestation³². This second mesodermal theory may also explain the presence of congenital skin masses associated with a median cleft of the upper lip, as a result of compensatory proliferation of mesodermal tissues, where the fusion of the facial processes occurs at the same embryonic stage³³, as well as the occurrence of pericallosal lipoma, as a result of mesodermal inclusion within the closing neural tube³⁴ [Au?4]. Several additional pathophysiological processes seem to be responsible for the development of pericallosal lipoma, including the persistence and abnormal differentiation of primitive meninx, which is a mesenchymal derivative of neural crests, into mature adipose tissue³⁵.

Thus, based on the literature review, a new definition for Pai syndrome spectrum is proposed: (1) a congenital nasal or mediofrontal skin mass or [Au?6] a mid-anterior alveolar process polyp (regardless of the pathological diagnosis, i.e., hamartoma, dermoid cyst, or lipoma) as an obligatory criterion, and at least one of the following criteria: (2) midline cleft of the upper lip and/or midline alveolar cleft, and/or (3) a pericallosal lipoma or [Au?6] interhemispheric lipoma in the case of corpus callosum dysgenesis (Fig. 4). Of the 60 cases described in the literature, 55 meet these proposed diagnostic criteria. The remaining five patients presented with only one criterion; however, an MRI was not performed in two of these patients [Au?4].

[Figure 4 here]

Perspectives: establishing differential diagnoses and elucidating the aetiology of Pai syndrome

The molecular basis of Pai syndrome remains unknown. To date, only one de novo reciprocal translocation, 46,X,t(X;16)(q28;q11.2), has been reported in a case of Pai syndrome including a midline cleft of the upper lip, pedunculated skin masses of the nasal septum, frontal bossing, hypertelorism, short and down-slanting palpebral fissures, short stature, and mental retardation²⁵. This phenotype overlaps with frontonasal dysplasia, as observed in 32% of the cases of Pai syndrome described in the literature review and in one patient in the case series. In patients presenting with Pai syndrome and additional features, chromosomal abnormalities may be detected more frequently, especially when the phenotype overlaps with those observed in frontonasal dysplasia spectrum³⁶ or in Goldenhar/OAV spectrum and oculoauriculofrontonasal syndrome, which is associated with features of both frontonasal dysplasia and OAV spectrum. Establishing consensus diagnostic criteria for Pai syndrome and performing systematic genetic investigations, including clinical evaluation by a geneticist and comparative genomic hybridization array, are necessary to better delineate Pai syndrome spectrum and thus shed light on its molecular basis.

Conclusions

Through the analysis of the case series patients and a literature review, this study was able to further analyse the phenotype of Pai syndrome, leading to the proposed new diagnostic criteria, i.e., (1) a congenital nasal or mediofrontal skin mass or [Au?6] a mid-anterior alveolar process polyp (regardless of the pathological diagnosis, i.e., hamartoma, dermoid cyst, or lipoma) as an obligatory criterion, and at least one of the following criteria: (2) midline cleft of the upper lip and/or midline alveolar cleft,

and/or (3) a pericallosal lipoma or [Au?6] interhemispheric lipoma in the case of corpus callosum dysgenesis.

Systematic cerebral imaging is recommended in the event that midline facial skin masses and/or a midline cleft lip is encountered, as well as ophthalmological assessment and evaluation by a geneticist to detect additional abnormalities and to establish the differential diagnosis of Pai syndrome or overlapping syndromes [Au?4]. This may help to better delineate the spectrum of Pai syndrome and elucidate its aetiology.

Funding

No funding was secured for this study.

Competing interests

All authors have no conflict of interest to declare.

Ethical approval

As a retrospective case series study, ethics committee approval was not required by the study institution [Au?4].

Patient consent

Informed consent was collected from the patients' parents for the publication of the photographs prior to submission.

References

1. Wu E, Vargevik K, Slavotinek AM. Subtypes of frontonasal dysplasia are useful in determining clinical prognosis. *Am J Med Genet A* 2007; 143A: 3069–3078. doi: 10.1002/ajmg.a.31963
2. Guion-Almeida ML, Mellado C, Beltrán C, Richieri-Costa A. Pai syndrome: report of seven South American patients. *Am J Med Genet A* 2007; 143A: 3273–3279. doi: 10.1002/ajmg.a.31962
3. Lederer D, Wilson B, Lefevre P, Poorten VV, Kirkham N, Mitra D, Verellen-Dumoulin C, Devriendt K. Atypical findings in three patients with Pai syndrome and literature review. *Am J Med Genet A* 2012; 158A: 2899–2904. doi: 10.1002/ajmg.a.35592
4. Pai GS, Levkoff AH, Leithiser RE, Reynolds JF. Median cleft of the upper lip associated with lipomas of the central nervous system and cutaneous polyps. *Am J Med Genet* 1987; 26: 921–924. doi: 10.1002/ajmg.1320260421
5. Castori M, Rinaldi R, Bianchi A, Caponetti A, Assumma M, Grammatico P. Pai syndrome: first patient with agenesis of the corpus callosum and literature review. *Birth Defects Res A Clin Mol Teratol* 2007; 79: 673–679. doi: 10.1002/bdra.20392
6. Lees MM, Connelly F, Kangesu L, Sommerlad B, Barnicoat A. Midline cleft lip and nasal dermoids over five generations: a distinct entity or autosomal dominant Pai syndrome? *Clin Dysmorphol* 2006; 15: 155–159. doi: 10.1097/01.mcd.0000204987.61990.f0
7. Vaccarella F, Pini Prato A, Fasciolo A, Pisano M, Carlini C, Seymandi PL. Phenotypic variability of Pai syndrome: report of two patients and review of the

literature. *Int J Oral Maxillofac Surg* 2008; 37: 1059–1064. doi:

10.1016/j.ijom.2008.06.007

8. Mishima K, Mori Y, Minami K, Sakuda M, Sugahara T. A case of Pai

syndrome. *Plast Reconstr Surg* 1999; 103: 166–170. doi: 10.1097/00006534-

199901000-00026

9. Szeto C, Tewfik TL, Jewer D, Rideout A. Pai syndrome (median cleft palate,

cutaneous nasal polyp, and midline lipoma of the corpus callosum): a case report and

literature review. *Int J Pediatr Otorhinolaryngol* 2005; 69: 1247–1252. doi:

10.1016/j.ijporl.2005.01.038

10. Imai Y, Seino H, Toriyabe S, Tachi M, Yamada A. Cases of true and false

median cleft with polypoid masses in the facial midline. *Cleft Palate Craniofac J*

2007; 44: 667–672. doi: 10.1597/06-120.1

11. Huckstadt V, Heis Mendoza ME, Moresco A, Obregon MG. [Pai syndrome:

two new cases with unusual manifestations]. *Arch Argent Pediatr* 2018; 116: e336–

e340. doi: 10.5546/aap.2018.e336

12. Melloni-Magnelli LF, de la Garza-Giacomán R, Martínez-Leija H, Guzmán-

Rodríguez R. Primer caso clínico de Síndrome de Pai en México. *Cirugía Plástica*

Ibero-Latinoamericana 2015; 41: 183–189. doi: 10.4321/S0376-78922015000200010

13. Li E, Galvin JA. Ophthalmic abnormalities of Pai syndrome: a case report and

review of literature. *Ophthalmic Genet* 2017; 14: 1–4. doi:

10.1080/13816810.2017.1381979

14. Iregbulem LM. Midline clefts of the upper lip. *Br J Plast Surg* 1978; 31: 63–

65. doi: 10.1016/0007-1226(78)90018-8

15. Preece JM, Kearns DB, Wickersham JK, Grace AR, Bailey CM. Nasal lipoma.

J Laryngol Otol 2007; 102: 1044–1046. doi: 10.1017/S0022215100107224

16. Morgan DW, Evans JN. Developmental nasal anomalies. *J Laryngol Otol* 2007; 104: 394–403. doi: 10.1017/S0022215100158542
17. Hollis LJ, Bailey CM, Albert DM, Hosni A. Nasal lipomas presenting as part of a syndromic diagnosis. *J Laryngol Otol* 2007; 110: 269–271. doi: 10.1017/S0022215100133389
18. Patil SB, Harsh S. Lipoma of columella with septal extension in Pai syndrome: report of a rare case. *BMC Ear Nose Throat Disord* 2017; 17: 657. doi: 10.1186/s12901-017-0035-y
19. AbdollahiFakhim S, Bayazian G, Notash R. Nasal septal lipoma in a child: Pai syndrome or not? *Int J Pediatr Otorhinolaryngol* 2014; 78: 697–700. doi: 10.1016/j.ijporl.2014.01.027
20. Savasta S, Chiapedi S, Perrini S, Tognato E, Corsano L, Chiara A. Pai syndrome: a further report of a case with bifid nose, lipoma, and agenesis of the corpus callosum. *Child Nerv Syst* 2008; 24: 773–776. doi: 10.1007/s00381-008-0613-9
21. Abdelmaaboud M, Nimeri N. Pai syndrome: first reported case in Qatar and review of literature of previously published cases. *BMJ Case Rep* 2012; 2012: bcr0220125940–0. doi: 10.1136/bcr-02-2012-5940
22. Chousta A, Ville D, James I, Foray P, Bisch C, Depardon P, Rudigoz RC, Guibaud L. Pericallosal lipoma associated with Pai syndrome: prenatal imaging findings. *Ultrasound Obstet Gynecol* 2008; 32: 708–710. doi: 10.1002/uog.6150
23. Yildiz H, Hakyemez B, Koroglu M, Yesildag A, Baykal B. Intracranial lipomas: importance of localization. *Neuroradiology* 2005; 48: 1–7. doi: 10.1007/s00234-005-0001-z

24. Truwit CL, Barkovich AJ. Pathogenesis of intracranial lipoma: an MR study in 42 patients. *Am J Roentgenol* 1990; 155: 855–864. doi: 10.2214/ajr.155.4.2119122
25. Masuno M, Imaizumi K, Fukushima Y, Tanaka Y, Ishii T, Nakamura M, Kuroki Y. Median cleft of upper lip and pedunculated skin masses associated with de novo reciprocal translocation 46,X,t(X;16)(q28;q11.2). *J Med Genet* 1997; 34: 952–954. doi: 10.1136/jmg.34.11.952
26. Guion-Almeida ML, Richieri-Costa A, Saavedra D, Cohen MM Jr. Frontonasal dysplasia: analysis of 21 cases and literature review. *Int J Oral Maxillofac Surg* 1996; 25: 91–97. doi: 10.1016/S0901-5027(96)80048-8
27. Azurdia J, Burke L, Laub DJ. Pai syndrome: median cleft lip, corpus callosum lipoma, and fibroepithelial skin tag. *Eplasty* 2014; 14: ic7. eCollection 2014.
28. Guion-Almeida ML, Richieri-Costa A. Frontonasal dysplasia, severe neuropsychological delay, and midline central nervous system anomalies: report of 10 Brazilian male patients. *Am J Med Genet A* 2009; 149A: 1006–1011. doi: 10.1002/ajmg.a.32717
29. Kieslich M, Ehlers S, Bollinger M, Jacobi G. Midline developmental anomalies with lipomas in the corpus callosum region. *J Child Neurol* 2016; 15: 85–89. doi: 10.1177/088307380001500205
30. Ocak Z, Yazicioglu HF, Aygun M, Ilter MK, Ozlu T. Prenatal detection of Pai syndrome without cleft lip and palate: a case report. *Genet Couns* 2013; 24: 1–5.
31. Stark RB. The pathogenesis of harelip and cleft palate. *Plast Reconstr Surg* 1954; 13: 20–39. doi: 10.1097/00006534-195401000-00003
32. Veau V. *Bec-de-lièvre*. Paris: Masson, 1938.

33. Nakamura J, Tomonari H, Goto S. True median cleft of the upper lip associated with three pedunculated club- shaped skin masses. *Plast Reconstr Surg* 1985; 75: 727–731. doi: 10.1097/00006534-198505000-00022
34. Wilberger JE Jr, Maroon JC, Prostko ER, Baghai P, Beckman I, Deeb Z. Magnetic resonance imaging and intraoperative neurosonography in syringomyelia. *Neurosurgery* 1987; 20: 599–605. doi: 10.1097/00006123-198704000-00016
35. Truwit CL, Barkovich AJ. Pathogenesis of intracranial lipoma: an MR study in 42 patients. *Am J Roentgenol* 1990; 155: 855–864. doi: 10.2214/ajr.155.4.2119122
36. Ribeiro-Bicudo LA, Quiezi RG, Guion-Almeida ML, Legnaro C, Richieri-Costa A. Exclusion of mutations in TGIF, ALX3, and ALX4 genes in patients with the syndrome of frontonasal dysgenesis, callosal agenesis, basal encephalocele, and eye anomalies. *Am J Med Genet A* 2012; 158A: 1233–1235. doi: 10.1002/ajmg.a.35305

Figure captions

Fig. 1. Frontal views of the patients in this series. (A) Patient 1 at the age of 1 year, showing the midline cleft lip and polypoid mass attached to the right side of the nasal septum. (B) Patient 2 at the age of 8 days, showing a large midline cleft lip and a pedunculated skin mass attached to the columella. (C) Patient 3 at the age of 5 months, showing a midline cleft lip, a bifid nose, hypertelorism, wide forehead, down-slanting palpebral fissures, and a pedunculated skin mass attached to the right nostril. (D) Patient 5 at the age of 1 month, showing a midline cleft lip, a skin mass of the alveolar cleft arising in the midline lip cleft, and a pedunculated skin mass of the right nostril (pedicled on the nasal septum). (E) Patient 6 at the age of 1 month, showing a midline cleft lip, a skin mass of the alveolar cleft arising in the midline lip cleft, and a pedunculated skin mass of the right nostril (pedicled on the nasal septum) [Au?7]. (F) Patient 7 at the age of 1 month, showing a right paramedian cleft lip, a median notch, and a pedunculated skin mass of the cleft lip, with extension into the right nostril. Note: Informed consent for publication of the frontal photograph was not obtained for patient 4.

Fig. 2. Additional facial features in patients 1, 2, and 4. (A) Intraoral view of patient 1: bifid labial frenum of the maxilla and midline alveolar cleft. (B) Lateral view of patient 2 at the age of 8 days, showing pre-auricular skin tags. (C) Frontal view of patient 4 at the age of 1 month, depicting a pedunculated skin mass attached to the nasal tip and a bifid frenum.

Fig. 3. Cerebral MRI depicting two types of pericallosal lipoma (sagittal plane): (A) tubulonodular-type lipoma of the corpus callosum (patient 1); (B) curvilinear-type lipoma of the corpus callosum (patient 4).

Fig. 4. Flowchart of new diagnostic criteria for Pai syndrome and suggested investigations. [Au?8]

A. Patient 1

B. Patient 2

C. Patient 3

D. Patient 5

E. Patient 6

F. Patient 7

A. Patient 1

B. Patient 2

C. Patient 4

A. Patient 1

B. Patient 4

Mandatory criteria:

A. ≥ 1 congenital nasal or mediofrontal and/or midanterior alveolar process skin mass

AND

B. Median cleft of the upper lip and/or median alveolar cleft

AND/OR

C. Pericallosal lipoma and/or interhemispheric lipoma in case of corpus callosum dysgenesis

Systematic cerebral imaging in case of A or B or A + B

Positive diagnosis of « Pai syndrome » if A + B, A + C, or A + B + C

Detailed physical examination including complete ophthalmological screening

« Isolated » Pai syndrome
if no other abnormalities*

« Associated » Pai syndrome in case of
additional unrecognizable or
recognizable syndrome

**Differential
diagnoses**

* The definition of additional abnormalities exclude additional congenital facial skin masses of the midline structure

Table 1. Clinical, radiological, and pathological characteristics of the patients diagnosed with Pai syndrome included in the current series.

Patient	Sex	Cutaneous facial polyps	Orofacial cleft	Pericallosal lipoma	Additional findings	Genetic
Birth term (weeks)		(localization, number, and				investigations
Birth weight (g)		histopathological diagnosis)				
Birth height (cm)						
Patient 1	Female	Polypoid mass attached to the	Midline cleft of the	Tubulonodular	Bifid frenum of the upper	Not performed
37 weeks + 5 days		right nasal septum (1)	upper lip ^a	pericallosal lipoma	lip	
3070 g		(hamartoma)	(microform)			
48.5 cm		Nasal skin lesion of the nasal tip	Median alveolar			
(Figs. 1–3)		(1) (dermoid cyst)	cleft of premaxilla			
Patient 2	Female	Diverticula of the nasal dorsum	Complete midline	Tubulonodular	Bilateral ear deformities	46 XX
40 weeks		(1) (hamartoma)	cleft lip and palate	pericallosal lipoma	Agenesis of the olfactive	Comparative
2640 g		Pedunculated skin mass pedicled		Short corpus	[Au?10] bulb	genomic
47.5 cm		on columella (1) (hamartoma)		callosum	Left ocular dermoid	hybridization array:
(Figs. 1, 2)		Diverticula of the upper gingiva				normal
		(1) (hamartoma)				(holoprosencephaly

Three left pre-auricular
pedunculated [Au?9] masses
(one fibrochondroma and two
lipofibromas)

excluded)

Patient 3	Female	Skin mass of the right nostril (1)	Complete midline	Tubulonodular	Hypertelorism	Not performed
38 weeks		(hamartoma)	cleft of the upper lip	pericallosal lipoma	Bifid nose	
2850 g				Partial agenesis of	Median fistula	
48 cm				the corpus	(columella)	
(Fig. 1)				callosum	Down-slanting palpebral fissures	
Patient 4	Female	Pedunculated skin mass on the	Midline cleft of the	Curvilinear lipoma	Bifid frenum of the upper	46 XX
41 weeks + 5 days		nasal tip (1) (hamartoma)	upper lip ^a	of corpus callosum	lip	
2980 g			(microform)		High arched palate	
49 cm			Median alveolar		Small left choroid plexus	
(Figs. 2, 3)			cleft of premaxilla		cyst	
					Hyperpigmented spot in	

					the left iris	
Patient 5	Male	Pedunculated skin mass of the	Incomplete midline	None	None	Not performed
38 weeks		right nostril (pedicled on the	cleft of the upper lip			
2950 g		nasal septum) (1) (hamartoma)	and alveolus			
48 cm		Skin mass of the alveolar cleft				
(Fig. 1)		arising in the midline cleft lip				
		(hamartoma) (1)				
Patient 6	Female	Nasal skin mass (columella	None	Tubulonodular	None	Exome sequencing:
40 weeks		extending to the right alar dome)		pericallosal lipoma		no abnormalities
2950 g		(dermoid cyst) (1)				
48 cm						
(Fig. 1)						
Patient 7	Male	Pedunculated skin mass of the	Complete	None	Congenital stenosis of	Exome sequencing:
38 weeks		cleft lip, with extension into the	paramedian cleft lip		the right lacrimal duct	no abnormalities
2800 g		right nostril (1) (hamartoma)	and palate (right		Duplication of the	
48 cm			side)		vomerian [Au?11] bone	

(Fig. 1)

Midline alveolar
cleft

^aCharacterized by disrupted vermillion–cutaneous junction, hypoplastic mucosal free margin, and notching of the vermillion.